



EPA REG. NO. 67517-95-56
EPA EST. NO. 56-OH-1



ACTIVE INGREDIENT:
Bromethalin (CAS No. 63333-35-7) ... 0.01%
OTHER INGREDIENTS ... 99.99%
TOTAL 100.00%

Contains: One Bait Station with 2 x .5 oz. of
BAIT BLOCKS® and 30 x .5 oz. of Replacement
BAIT BLOCKS® • Net Wt. 1 lb. (16 oz.)

KEEP OUT OF REACH OF CHILDREN

CAUTION
SEE SIDE/BACK PANELS FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

ITEM #939



**MAY BE HARMFUL IF EATEN
BY CHILDREN OR PETS**

**FOR USE INDOORS AND OUTDOORS
MOISTURE RESISTANT BAIT**

House Mice usually consume a lethal dose in a single feeding,
but it may take two or more days for them to die.
Due to Stop Feed Action, house mice cease feeding after consuming a toxic dose.
This product is effective against anticoagulant-resistant house mice.

**REFILLABLE
MOUSE POISON
BAIT STATION**



ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals and birds. Dogs and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: THIS BAIT STATION MAY BE USED IN AND AROUND (WITHIN 50 FEET OF) BUILDINGS ACCESSIBLE TO CHILDREN AND PETS, CONSISTENT WITH USE RESTRICTIONS AND OTHER REQUIREMENTS INDICATED IN THIS LABEL.

To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Dispose of product container and unused, spoiled and unconsumed bait as specified on this label.

USE SITES: For indoor and outdoor use. This product may only be used to control house mice in and around homes and residential buildings.

USE RESTRICTIONS. Do not use this product in sewers. Do not place this ready-to-use bait station in any area where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not apply further than 50 feet from home or building. Do not deliberately remove bait from bait station for broadcasting. Do not sell this product in individual packages holding more than 1 pound of bait.

SELECTION OF TREATMENT AREAS: Determine areas where house mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in corners, burrows and concealed places, between floors and walls, or in locations where house mice or their signs have been seen. Remove as much alternative food accessible to house mice as possible.

APPLICATION DIRECTIONS: The active ingredient in this product is different from anticoagulant rodenticides. Individual house mice cease feeding after consuming a toxic dose.

House Mice: Place bait stations 8 to 12 feet apart in infested area. Highest bait consumption is expected to occur on the first day or two after treatment. After several days, inspect bait placements and replace with new bait at sites where there is heavy feeding or where there is continued evidence of house mouse activity. Continue baiting until all signs of house mouse activity have ceased.

While wearing gloves, collect and properly dispose of all dead house mice and used bait stations. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment.

PESTICIDE STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container, except as described in the directions for use.

If Empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not get in eyes, on skin, or on clothing. Keep away from children, domestic animals and pets. Any person who retrieves carcasses or used bait following application of this product must wear waterproof gloves.

USER SAFETY REQUIREMENTS:

As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID

Have product container or label with you when calling a Poison Control Center or doctor for treatment. You may also contact the Hot Line Number at 1-800-345-4735 for health concerns, emergency medical treatment information or pesticide incidents.

IF SWALLOWED

- Call a Poison Control Center, doctor, or HOT LINE NUMBER 1-800-345-4735 immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a Poison Control Center or doctor.
- Do not give anything by mouth to an unconscious person.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye.
- Call a Poison Control Center or doctor for treatment advice.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes
- Call a Poison Control Center, doctor, or HOT LINE NUMBER 1-800-345-4735 immediately for treatment advice.

TREATMENT FOR PET POISONING

If an animal eats this bait, call a veterinarian or 1-800-345-4735 at once.

NOTE TO PHYSICIAN OR VETERINARIAN


Contains the nerve poison bromethalin. This product is not an anticoagulant type rodenticide. For humans or animals ingesting bait and/or showing poisoning signs (such as muscle tremors, loss of hind limb use, or seizures for animals), limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.

INSTRUCTIONS FOR OPENING, REFILLING & RECLOSING THE MOUSE BAIT STATION

1. To open the bait station, put on gloves and insert the plastic key into the lock on top of the Bait Station lid. Push the key all the way down into the lock and turn key in the direction of the arrow to open the station lid.

2. REMOVE AND SAVE KEY

3. Remove the plastic bait holder and discard any remaining bait from both the holder and the Bait Station. Place one refill block of bait into each side of the plastic bait holder, with center holes facing upward.



4. Insert plastic bait holder with two refill baits into the **MIDDLE COMPARTMENT** of the Bait Station as shown below.

5. WITH KEY STILL REMOVED and plastic bait holder and refills in place, snap the lid of the Bait Station shut by pressing down on the lock.

6. CLEAN AROUND THE BAIT STATION AREA BY REMOVING ALL REMAINING BAIT & BAIT CRUMBS. ENSURE THAT GLOVES AND BAIT ARE DISPOSED OUT OF REACH OF CHILDREN, PETS AND WILDLIFE.



JT EATON™ is a trademark and BAIT BLOCKS® and ANSWER® are registered trademarks of the J.T. EATON & CO., INC., Twinsburg, OH 44087 • U.S.A.
BITREX® is a registered trademark of Macfarlan Smith, Scotland ©2012 J.T. EATON & CO., INC.

EPA01132011-INV/011112A FNNW939016JTA

J.T. EATON & CO., INC.,
1393 East Highland Road
Twinsburg, OH 44087 • U.S.A. •
www.jteaton.com

